

ANTIBODIES AGAINST HUMAN IL-12

5
elms
A1
ens 92

~~This is a continuation application of Serial No. 09/232,522, filed January 19, 1999,~~
~~which claims priority under 35 USC § 119(e) of prior Provisional Application Serial No. 60/072,333,~~
~~filed January 23, 1998.~~

FIELD OF THE INVENTION

This invention relates generally to IL-12 antibodies, and more specifically to anti-human IL-12 polyclonal and monoclonal antibodies.

BACKGROUND OF THE INVENTION

Interleukin-12 (IL-12), formerly known as cytotoxic lymphocyte maturation factor or natural killer cell stimulatory factor, is a 75-kDa (p75) heterodimeric cytokine composed of disulfide bonded 40-kDa (p40) and 35-kDa (p35) subunits. The p40 and p35 subunits are polypeptides which contain 306 amino acid residues and 197 amino acid residues, respectively (Gubler U., et al., Proc. Natl. Acad. Sci. USA, Vol. 88, 4143-4147 (1991)).

The p75 heterodimer is the biologically active form of IL-12 (Gubler, U., et al., 1991, Proc. Natl. Acad. Sci. USA, 88: 4143; Wolf, S.F., et al., 1991, J. Immunol., 146: 3074). The IL-12 p75 heterodimer both activates and boosts cell mediated immune responses against foreign antigens by stimulating production of Th1 helper cells, stimulating activated T and natural killer (NK) cells,

enhancing lytic activity of NK/LAK cells, and stimulating production of IFN- γ by resting and activated T and NK cells.

5 The p40 subunit of IL-12 has been shown to be produced in excess of the p35 subunit and is found in both monomeric and dimeric forms (Podlaski, F.J., et al., 1992, Arch. Biochem. Biophys. 294: 230; D'Andrea, A., et al., 1992, J. Exp. Med., 176: 1387). IL-12 p40 homodimer is a potent IL-12 antagonist (Ling, P., et al., 1995, J. Immunol., 154: 116; Gillessen, S., et al., 1995, Eur. J. Immunol., 25: 200). In contrast to the p40 subunit, the p35 subunit of IL-12 has no known biological activity, and the p35 protein has only been found in association with the p40 subunit as part of the IL-12 p75 heterodimer. Therefore, there are two important types of epitopes presented by human IL-12: (1) epitopes presented by the p40 subunit; and (2) epitopes presented by the three dimensional conformation of the IL-12 p75 heterodimer. Consequently, we designate antibodies that recognize epitopes present on the IL-12 p75 heterodimeric protein but do not recognize epitopes present on the IL-12 p40 subunit protein "heterodimer specific" antibodies.

It has been found that known IL-12 antibodies are not optimally effective in substantially neutralizing IL-12 bioactivity. IL-12 antibodies which immunologically react with the p40 subunit do not optimally block the bioactivity of human IL-12. For example, use of antibodies which react with epitopes presented by the p40 subunit is particularly problematic because production of IL-12 p75 heterodimer has been shown to result in excess inactive p40 subunits relative to bioactive p75 heterodimer (Podlaski, F.J., 1992, Arch. Biochem. Biophys. 294: 230; D'Andrea, A., et al., 1992, J. Exp. Med., 176: 1387). As a result, the p40 antibodies are not as effective as heterodimer specific antibodies in reducing detrimental effects of IL-12 because the p40 subunit alone is not bioactive, and p40 antibodies tend to bind to the inactive p40 subunits rather than those p40 subunits that are part of a bioactive p75 heterodimer.

Even known antibodies which react only with the p75 heterodimer, do not effectively neutralize IL-12 bioactivity. For example, a previously identified IL-12 p75 heterodimer specific antibody, called 20C2 (Chizzonite et al., Cytokine, 6: A82a (1994) and D'Andrea et al., J. Exp. Med., Vol. 176, 1387-1398 (1992)), cannot substantially block human IL-12 stimulated PHA-activated lymphoblast proliferation and IFN- γ production.

Heterodimer specific antibodies which more effectively neutralize IL-12 bioactivity are needed to reduce detrimental effects of IL-12. Increased levels of IL-12 in serum or tissue are known to be involved in the development and progression of autoimmune disorders. Thus, IL-12 antibodies are useful antagonists for controlling diseases with pathologies that are mediated through immune mechanisms, particularly, diseases associated with aberrant Th1-type helper cell activity. Examples of such autoimmune disorders include multiple sclerosis, inflammatory bowel disease (IBD) including Crohn's disease and ulcerative colitis, rheumatoid arthritis and autoimmune diabetes mellitus. Other disease conditions which have been shown to benefit from the administration of IL-12 antibodies include transplantation/graft-versus-host disease and septic shock.

In accordance with this invention, it has been found that IL-12 antibodies obtained from a mammal deficient in the gene encoding the p35 subunit and/or the gene encoding the p40 subunit substantially neutralize IL-12 bioactivity.

SUMMARY OF THE INVENTION

In accordance with this invention, for the first time, antibodies which substantially neutralize the bioactivity of human IL-12 are produced using the methods described herein. Unlike other IL-12 p75 heterodimer specific antibodies, the heterodimer specific antibodies of the present invention

neutralize at least 90% of human IL-12 bioactivity. In addition, IL-12 p75 heterodimer specific antibodies of the present invention cross react with rhesus monkey IL-12.

The p75 heterodimer specific IL-12 antibodies described herein are effective therapeutic agents for use in blocking IL-12 bioactivity to treat conditions mediated by undesirable IL-12 stimulated immunological responses. The highly neutralizing heterodimer specific IL-12 antibodies described herein are particularly useful inhibitors of IL-12 stimulated PHA-activated human lymphoblast proliferation and IFN- γ production by PHA-activated human lymphoblasts.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a graph showing capture of ^{125}I -labeled human IL-12 by antibodies contained in supernatants from hybridomas HIL-12F3-5F2 (herein referred to as "5F2"), HIL-12F3-16F2 (herein referred to as "16F2"), HIL-12F3-16G2 (herein referred to as "16G2"), HIL-12F3-20E11 (herein referred to as "20E11"), and HIL-12F1-17E2 (herein referred to as "17E2") (open bars). The presence of unlabeled human IL-12 p40 subunit during the immunoprecipitation reaction (solid bars) did not block capture of ^{125}I -labeled human IL-12 by monoclonal antibodies 5F2, 16F2, 16G2 and 20E11, demonstrating that these antibodies do not have high affinity for the IL-12 p40 subunit alone.

Fig. 2 shows isoelectric focusing patterns of p75 heterodimer specific anti-human IL-12 monoclonal antibodies 20C2, 16G2, 16F2, 20E11, and 5F2. As shown in Fig. 2, monoclonal antibodies 20C2, 20E11 and 5F2 are unique immunoglobulins. Monoclonal antibodies 16G2 and 16F2 appear identical by isoelectric focusing, but both are different from 20C2, 20E11 and 5F2.

Fig. 3 is a plot showing inhibition of natural human IL-12 stimulated PHA-activated human lymphoblast proliferation by p75 heterodimer specific IL-12 monoclonal antibodies 20C2 (+), 16G2 (Δ), 16F2 (O), 20E11 (:), and 5F2 (s). Inhibition of natural human IL-12 stimulated PHA-activated

human lymphoblast proliferation was determined with respect to the level of 0.25 ng/ml human IL-12 stimulated PHA-activated human lymphoblast proliferation in the absence of IL-12 antibodies, shown in Fig. 3 as a horizontal dotted line at 9940 cpm, and background levels of PHA-activated human lymphoblast proliferation, i.e., in the absence of both IL-12 and IL-12 antibodies, shown in Fig. 3 as a horizontal dotted line at 1480 cpm. As shown in Fig. 3, IL-12 monoclonal antibodies, 16G2 (Δ), 16F2 (\circ), 20E11 ($:$) and 5F2 (s) inhibit 0.25 ng/ml human IL-12 stimulated PHA-activated lymphoblast proliferation by at least 90%. In contrast, as shown in Fig. 3, previously known 20C2 (+) antibody does not substantially inhibit IL-12 stimulated PHA-activated human lymphoblast proliferation.

Fig. 4 is a plot showing inhibition of rhesus monkey IL-12 stimulated PHA-activated human lymphoblast proliferation by p75 heterodimer specific IL-12 monoclonal antibodies 16G2 (Δ), 16F2 (\circ), 20E11 ($:$) and 5F2 (s) of the present invention compared to the previously known 20C2 (+) antibody. The level of lymphoblast proliferation in the presence of 0.5 ng/ml rhesus monkey IL-12 and in the absence of IL-12 antibodies is represented by the horizontal dotted line at the upper end of the plot. The background level of lymphoblast proliferation, i.e., in the absence of both IL-12 and IL-12 antibodies is represented by a horizontal dotted line at the lower end of the plot. As shown in Fig. 4, the antibodies of the present invention are potent inhibitors of rhesus monkey IL-12 stimulated PHA-activated lymphoblast proliferation, in contrast to the 20C2 (+) antibody which has a minimal inhibitory effect on rhesus monkey IL-12 stimulated lymphoblast proliferation.

Fig. 5 is a plot showing inhibition of IFN- γ production by p75 heterodimer specific monoclonal antibodies, 16F2 (\circ), 16G2 (\blacksquare), 20E11 (s), 5F2 (\bullet) and 20C2 (Q). As shown in Fig. 5, antibodies 16F2 (\circ), 16G2 (\blacksquare), 20E11 (s) and 5F2 (\bullet) inhibit 0.25 ng/ml human IL-12 stimulated IFN- γ production by at least 90%. The dashed horizontal line at the lower end of the plot represents background IFN- γ production in the absence of IL-12. In contrast, as shown in Fig. 5, the 20C2 (Q)

monoclonal antibody is unable to inhibit 0.25 ng/ml IL-12 stimulated IFN- γ production by more than 65%.

1NSC1

5 Fig. 6 is a nucleotide sequence encoding a portion of the heavy chain variable region of the p75 heterodimer specific 16G2 antibody, and the amino acid sequence deduced from this nucleotide sequence.

1NSC2

10 Fig. 7 is a nucleotide sequence encoding a portion of the heavy chain variable region of the p75 heterodimer specific 20E11 antibody, and the amino acid sequence deduced from this nucleotide sequence.

DETAILED DESCRIPTION OF THE INVENTION

15 In accordance with the present invention, it has been found that when IL-12 antibodies are produced from mammals deficient in the gene encoding the p35 IL-12 subunit and/or the gene encoding the p40 IL-12 subunit, IL-12 antibodies are obtained which selectively immunologically react with epitopes of the p75 heterodimer of IL-12, and are identified by their ability to selectively immunologically react with the p75 heterodimer of human IL-12, but not immunologically react with the p40 subunit alone.

20 Unlike previously known IL-12 p75 antibodies, antibodies which substantially neutralize the bioactivity of human IL-12, i.e., neutralize at least about 90% bioactivity of human IL-12, are produced by the methods described herein. In addition, IL-12 p75 heterodimer specific antibodies of the present invention cross react with rhesus monkey IL-12.

25 The IL-12 antibodies described herein neutralize at least about 90% bioactivity of human IL-12 by inhibiting at least about 90% IL-12 induced PHA-activated human lymphoblast proliferation at

concentrations of at least about 0.5 µg/ml, and/or inhibiting at least about 90% of IL-12 stimulated IFN-γ production by PHA-activated human lymphoblasts at concentrations of at least about 0.5 µg/ml. Furthermore, the antibodies described herein have been shown to specifically inhibit IL-12-induced, but not IL-2-induced, proliferation of PHA-activated human lymphoblasts. PHA-activated lymphoblasts are prepared as follows. Peripheral blood mononuclear cells (PBMC) were isolated (Gately et al., J. Natl. Cancer Inst., 69:1245 (1982)) and stimulated with 0.1% PHA-P (Difco Labs., Detroit, MI). After 3 days, the cultures were split 1:1 with fresh medium and recombinant 50 U/ml human IL-2 (provided by Dr. F. Khan, Hoffmann-La Roche Inc.) as described in Gately, M.K., Chizzonite, R. and Presky, D.H., Measurement of human and mouse interleukin 12, *Current Protocols in Immunology*, vol. 1. J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, and W. Strober, eds., John Wiley & Sons, Inc., New York, 1995, pp. 6.16.1-6.16.15. The PHA-activated lymphoblasts were used after an additional one day incubation period.

In accordance with the present invention, the IL-12 antibodies are identified for their ability to selectively bind the epitope presented by the p75 heterodimer, but not immunologically react with any epitope presented by the p40 subunit. This selectivity is defined by the fact that the IL-12 antibodies of this invention will react, at a certain minimal concentration, with an epitope solely presented by a given amount of the p75 heterodimer but will not react at that concentration with an epitope presented by the p40 subunit of that same given amount of this p75 heterodimer. In this way the antibodies of this invention have a higher affinity for an epitope solely presented by the p75 heterodimer than any epitope presented by the p40 subunit. Any conventional assay for identifying selective binding of the antibodies to the p75 heterodimer can be used. Generally, in such an assay, antibodies are incubated with human IL-12 p75 heterodimer to determine if the antibodies bind the p75 heterodimer. The antibodies are also incubated with human IL-12 p75 heterodimer in the presence and absence of the p40 subunit to determine if the presence of the p40 subunit blocks antibody binding or capture of the p75 heterodimer. For example, competitive immunoprecipitation assays (see Example 7 herein) can be

used to demonstrate that the antibodies described herein selectively immunologically react with the p75 heterodimer of human IL-12, but are not immunologically reactive with the p40 subunit alone.

In accordance with the present invention, the IL-12 antibodies described herein are produced through the use of knock-out mammals. The knock-out mammals are deficient in the gene encoding the p35 subunit and/or the gene encoding the IL-12 p40 subunit and thus, do not express the IL-12 p75 heterodimer. When immunized with the IL-12 p75 heterodimer, the IL-12 p35 subunit deficient and/or the IL-12 p40 subunit deficient knock-out mammal recognizes the IL-12 p75 heterodimer as foreign and produces antibodies thereto. Preferably, the knock-out mammal is a mouse. In accordance with the present invention knock-out mammals are produced by methods that have been described in the art. Knock-out mammals can be produced by conventional means such as by mutation of the gene encoding the p35 IL-12 subunit and/or the p40 IL-12 subunit. For example, mice carrying a mutation in the IL-12 p35 subunit gene can be produced as described by Mattner, F., et al., *Eur. J. Immunol.*, 26:1553-1559 (1996). Mice carrying a mutation in the IL-12 p40 subunit gene can be produced as described by Magram, J., et al., *Immunity*, 4: 471-481 (1996).

In accordance with the present invention, polyclonal and monoclonal antibodies that selectively immunologically react with the p75 heterodimer of human IL-12 are produced from activated cells of the aforementioned knock-out mammal by any conventional means known in the art. Generally, the antibodies are produced by (a) immunizing a "knock-out" mammal deficient in a gene encoding the p35 subunit and/or the p40 subunit with human p75 heterodimer to produce antibodies; (b) obtaining antibodies from the immunized mammal; and (c) screening the antibodies for their ability to selectively bind the epitope presented by the p75 heterodimer to obtain the selectively binding antibodies.

The IL-12 monoclonal antibodies of the present invention which selectively immunologically react with the human IL-12 p75 heterodimer are generally produced by a method including the following steps:

(1) immunizing a knock-out mammal, such as, for example, a mouse deficient in the gene encoding the IL-12 p35 subunit and/or IL-12 p40 subunit, with human IL-12 p75 heterodimer;

(2) selecting cells from the immunized knock-out mammal that have been activated to express antibodies against IL-12, such as, splenocytes or lymph node cells;

5 (3) fusing the harvested cells to myeloma cells to form hybridoma cells;

(4) selecting hybridoma cells which secrete antibodies that recognize human IL-12, for example, by testing hybridoma conditioned medium for the presence of anti-human IL-12 antibodies, for example, through the use of ELISA or immunoprecipitation assays employing labeled or unlabeled human IL-12; and

10 (5) determining if the antibodies are p75 heterodimer specific by demonstrating that the antibodies immunologically react with an epitope of the p75 IL-12 heterodimer, but are not immunologically reactive with any epitope of the p40 subunit, by incubating the antibodies with human IL-12 p75 heterodimer to determine if the antibodies bind the p75 heterodimer, and then incubating the antibodies with human IL-12 p75 heterodimer in the presence and absence of the p40 subunit to determine if the presence of the p40 subunit blocks antibody binding or capture of the p75 heterodimer. For example, competitive immunoprecipitation assays (see Example 7 herein) can be used to demonstrate that the antibodies described herein selectively immunologically react with the p75 heterodimer of human IL-12, but are not immunologically reactive with the p40 subunit alone.

20 The method for producing the p75 heterodimer specific IL-12 monoclonal antibodies of the present invention can further comprise the step of determining the ability of the heterodimer specific IL-12 antibodies to inhibit both human and rhesus monkey IL-12 bioactivity in any *in vitro* or *in vivo* assay system for IL-12 bioactivity, such as assays for determining IL-12-stimulated proliferation of activated lymphocytes, IL-12-stimulated production of IFN- γ , or IL-12-stimulated enhancement of
25 cytolytic activity.

The anti-human IL-12 antibodies of the present invention can be isolated to substantially pure form by standard methods known in the art, such as, for example, ammonium sulfate precipitation, affinity chromatography, or ion exchange chromatography.

Variations of the method for obtaining the antibodies of the present invention are also encompassed within the present invention. Methods known in the art such as, for example, Western blotting, competitive immunoprecipitation assays, or cross-blocking binding assays can be used to determine if the antibodies are p75 heterodimer specific.

In addition to mice, mammals such as rats and rabbits deficient in the IL-12 p35 subunit gene and/or IL-12 p40 subunit gene, can be immunized with the IL-12 p75 heterodimer to produce the antibodies described herein. The deficiency or mutation in the IL-12 p35 subunit gene and/or IL-12 p40 subunit gene can be any deficiency or mutation that results in lack of expression of IL-12 p75 heterodimer. Furthermore, any conventional method for obtaining mammalian cells carrying a mutation in the IL-12 p35 subunit gene and/or IL-12 p40 subunit gene which results in IL-12 p75 deficient phenotype can be used.

In accordance with the present invention, activated mammalian cells expressing antibodies to the human IL-12 p75 heterodimer can be obtained by immunizing a mouse or other mammal with natural human IL-12 or recombinant IL-12. Natural human IL-12 and recombinant human IL-12 can be prepared by any conventional technique known in the art, such as the techniques provided in the examples herein.

Suitable myeloma cell lines, i.e., fusion partners, for use in producing the hybridomas that secrete the IL-12 antibodies of the present invention include myeloma cell lines well known in the art, such as, for example, SP 2/0 and NS/O cell lines. SP2/0 mouse myeloma cells are preferred. Preferably, the myeloma fusion partner and the mammalian cell activated against the IL-12 p75 heterodimer are derived from the same species.

Hybridoma cells producing the antibodies of the present invention can be selected and isolated by any conventional methods known in the art. Preferably, myeloma cells and lymphocytes activated against the IL-12 p75 heterodimer are cultured together in media containing a selection agent capable of killing the myeloma cells but not the lymphocytes. Hybridomas are formed from myeloma cells that fuse with the lymphocytes activated against the IL-12 p75 heterodimer. Such hybridoma cells are capable of growing in the media containing the selection agent because DNA of the lymphocytes supplies to the myeloma cell line the necessary gene encoding an enzyme that prevents the toxic effects of the selection agent by allowing an alternate metabolic pathway to replace the metabolic pathway blocked by the selection agent. Any unfused lymphocytes die because they are not transformed and have short, finite lifetimes *in vitro*. In accordance with the present invention a suitable selection agent for use in selecting out hybridoma cells is aminopterin. A preferred medium for cultivating the hybridoma cells is Iscove's Modified Dulbecco's Medium (IMDM) supplemented with 10% FBS (Hyclone), 100 Units/ml penicillin G (BioWhittaker), 100 µg/ml streptomycin (BioWhittaker), 250 ng/ml Fungizone (BioWhittaker), 2 mM glutamine (BioWhittaker), 100 µg/ml gentamicin sulfate (BioWhittaker), 50 µM 2-mercaptoethanol (BioRad), 100 µM hypoxanthine (Sigma), 400 nM aminopterin (Sigma), 16 µM thymidine (Sigma), and 2.5% P388D1 supernatant (produced as described by Nordan, R.P., et al., J. Immunol., 139:813 (1987)).

The potency of the IL-12 antibodies of the present invention is determined with respect to the concentration of IL-12 antibodies at which 50% of maximal inhibition of IL-12 bioactivity occurs as measured by IL-12-stimulated human lymphoblast proliferation or IFN-γ production assays. The anti-human IL-12 antibodies of the present invention exhibit higher potency than previously characterized heterodimer specific IL-12 antibodies. In addition, the anti-human antibodies of the present invention exhibit greater efficacy, as measured by the extent of maximal inhibition of IL-12-stimulated lymphocyte proliferation or IFN-γ production, than previously characterized heterodimer specific IL-12 antibodies.

The potency and efficacy of the antibodies described herein can be determined by any conventional assay known in the art, such as, for example, IL-12 induced lymphoblast proliferation assays or IFN- γ synthesis assays.

5

In accordance with the present invention, any conventional method known in the art can be used to determine inhibition of human IL-12 stimulated lymphoblast proliferation by the IL-12 antibodies. In general, human lymphocytes can be activated by a number of methods, including treatment with mitogenic lectins, *e.g.* phytohemagglutinin A (PHA), or other activating agents, alone or in combination, such as cytokines, phobol esters and ionophores, antibodies directed against cell surface molecules, or any other method which will lead to activation of the lymphocytes. The activated lymphocytes are then incubated with and without IL-12 in the absence or presence of the antibodies, and the rate of lymphocyte proliferation is measured by determining the rate of DNA synthesis by measuring the incorporation of ^3H -thymidine into DNA, by counting the number of cells present after various periods of treatment, or any other method that can be used to monitor the rate of cellular proliferation. Inhibition of proliferation is determined by comparing lymphocyte proliferation at a defined concentration of IL-12 in the absence and presence of various concentrations of anti-IL-12 antibodies.

20 In a standard lymphocyte proliferation assay, inhibition of human IL-12 stimulated PHA-activated human lymphoblast proliferation is determined with respect to levels of human IL-12 stimulated PHA-activated human lymphoblast proliferation without any added antibodies and background levels of PHA-activated human lymphoblast proliferation, *i.e.*, proliferation in the absence of both IL-12 and antibodies. In general, IL-12-stimulated levels of proliferation yield about 10,000-
25 80,000 cpm in our standard human lymphocyte proliferation assay, with background levels of proliferation yielding about 5,000-20,000 cpm. Due to the inherent variability between batches of stimulated PHA-activated human lymphoblasts, only assays in which the ratio of stimulated

proliferation to background proliferation (*i.e.* the stimulation index) was equal to or greater than 3 are considered valid for the measurement of IL-12-stimulated proliferation.

In accordance with the present invention, any conventional method for determining inhibition of IFN- γ production by the IL-12 antibodies can be used. For example, activated human lymphocytes, prepared as described herein, or activated human peripheral blood mononuclear cells (PBMC), prepared by treating whole blood or isolated PBMC with mitogenic agents including lectins, cytokines, phorbol esters, ionophores, or antibodies directed against cell surface molecules, alone or in combination, or by any other method which will lead to the production of activated human PBMC, are incubated with or without IL-12 and various other agents, *e.g.* IL-2 and/or IL-1 β , in the absence and presence of the antibodies. IFN- γ production is then determined, *e.g.* by sampling the culture medium and determining the concentration of IFN- γ by ELISA or any other method that can quantitatively measure IFN- γ . Inhibition of IFN- γ production is determined by comparing IFN- γ production at a defined concentration of IL-12 in the absence and presence of various concentrations of anti-IL-12 antibodies.

In a standard IFN- γ synthesis assay inhibition of IFN- γ is determined with respect to IL-12-stimulated IFN- γ production and background levels of IFN- γ production, *i.e.*, IFN- γ synthesis in the presence or absence of IL-12. In general, IL-12-stimulated levels of IFN- γ production are about 7-220 ng/ml, with background levels of production yielding about 1-3 ng/ml.

The antibodies herein neutralize rhesus monkey IL-12 bioactivity with a potency similar to their potency for inhibiting human IL-12 bioactivity, making them useful IL-12 antagonists for *in vivo* studies in the rhesus monkey. The increased potency and efficacy of these anti-human IL-12 antibodies and their cross reactivity with rhesus monkey IL-12 make them excellent candidates for designing effective IL-12 antagonists for use in humans.

INSC3)

In particular, the present invention provides four antibodies, 5F2, 16F2, 16G2 and 20E11 to the p75 heterodimer of human IL-12 having ATCC designation numbers HB-12446, HB-12447, HB-12449, and HB-12448, respectively. However, the present invention is not limited to these four antibodies. Any antibodies having the characteristics described herein are encompassed within the present invention.

INSC4)

Fig. 6 provides the nucleotide sequence encoding a portion of the heavy chain variable region of the p75 heterodimer specific 16G2 antibody and the amino acid sequence deduced from this nucleotide sequence. The nucleotide sequence encoding a portion of the heavy chain variable region of the p75 heterodimer specific 20E11 antibody and the amino acid sequence deduced from this nucleotide sequence is provided in Fig. 7. It will be understood by those skilled in the art that conservative amino acid changes can be made in the constant regions of the heterodimer specific IL-12 antibodies herein without significantly affecting the antigen binding specificity/affinity. Heterodimer specific IL-12 antibodies containing amino acid changes in the variable framework regions or complementary determining regions can be expected to have a greater effect on antigen binding specificity/affinity.

The IL-12 antibodies of the present invention can be complete antibodies including two full length heavy chains and two full length light chains. Alternatively, the IL-12 antibodies can be constructs such as single chain antibodies or "mini" antibodies that retain binding activity to one or more epitopes of the IL-12 p75 heterodimer. Such constructs can be prepared by methods known in the art such as, for example, the PCR mediated cloning and assembly of single chain antibodies for expression in E. coli (as described in Antibody Engineering, The practical approach series, J. McCafferty, H.R. Hoogenboom, and D. J. Chiswell, editors, Oxford University Press, 1996). In this type of construct, the variable portions of the heavy and light chains of an antibody molecule are PCR

amplified from cDNA. The resulting amplicons are then assembled, *e.g.* in a second PCR step, through a linker DNA that encodes a flexible protein linker composed of the amino acids GLY and SER. This linker allows the variable heavy and light chain portions to fold in such a way that the antigen binding pocket is regenerated and antigen is bound with affinities often comparable to the parent full-length dimeric immunoglobulin molecule.

The IL-12 antibodies of the present invention are useful antagonists for controlling diseases with pathologies that are mediated through immune mechanisms, particularly, diseases associated with increased IL-12 bioactivity that results in aberrant Th1-type helper cell activity. In accordance with the present invention, the IL-12 antibodies are used for treating autoimmune disorders in humans or other mammals, such as, for example, multiple sclerosis, rheumatoid arthritis, autoimmune diabetes mellitus, and inflammatory bowel disease (IBD) including Crohn's disease and ulcerative colitis. The antibodies described herein can also be used to treat other disease conditions which have been shown to benefit from the administration of IL-12 antibodies including, for example, transplantation/graft-versus-host disease and septic shock.

The dose ranges for the administration of the IL-12 antibodies herein may be determined by those of ordinary skill in the art without undue experimentation. In general, appropriate dosages are those which are large enough to produce the desired effect, *i.e.*, neutralizing at least 90% IL-12 bioactivity. However, the dosage should not be so large as to cause adverse side effects, such as unwanted cross-reactions, anaphylactic reactions, and the like. Generally, the dosage will vary with the age, condition, sex and extent of disease in the patient, counter indications, if any, immune tolerance and other such variables, to be adjusted by the individual physician.

The IL-12 antibodies may be administered parenterally by injection or by gradual perfusion over time. They can be administered intravenously, intramuscularly, or subcutaneously. Preparations

for parenteral administration include sterile or aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media.

5 Parenteral vehicles include sodium chloride solution, Ringers' dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers, such as those based on Ringer's dextrose, and the like. Preservatives and other additives may also be present, such as, for example, anti-microbials, anti-oxidants, chelating agents, inert gases and the like. See generally, *Remington's Pharmaceutical Science*, 16th Ed., Mack Eds., 1980, which is incorporated herein by reference.

Preferred dosages of the IL-12 antibodies of the present invention are from about 0.1 mg/kg to about 10 mg/kg, two to three times per week. However, the dosage and dosage schedule for administration of the IL-12 antibodies herein may vary depending on the individual to be treated, the antibody administered, and the variables discussed above. In accordance with the present invention, the IL-12 antibodies may be administered alone or in combination with other therapeutically active agents.

20 The anti-human IL-12 antibodies described herein may be humanized to form antibodies that possess the same or substantially similar affinity for the IL-12 p75 heterodimer as mammalian anti-human IL-12 antibodies, but are substantially non-immunogenic in humans. For example, a humanized IL-12 antibody in accordance with the present invention can include heavy and light chain framework regions of human antibodies. Preferably, the amino acid sequences of the humanized antibody framework regions are from about 60% to 95% identical to the donor framework regions. The
25 humanized antibodies may be produced by recombinant techniques well known in the art. Methods for producing humanized immunoglobulins are described in U.S. Patent No. 5,530,101, the disclosure of which is incorporated herein by reference.

EXAMPLE 1

Preparation of Natural Human IL-12

5

Blood was drawn from normal volunteer donors into syringes containing preservative-free heparin (Sigma, St. Louis, MO) to give a final concentration of ~ 5 units heparin/ml blood. One volume of heparinized blood was diluted into 9 volumes of medium consisting of a 1:1 mixture of RPMI 1640 and Dulbecco's modified Eagle's medium, supplemented with 0.1 mM nonessential amino acids, 60 µg/ml arginine HCl, 10 mM HEPES buffer, 2 mM L-glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin (all available from GIBCO BRL, Grand Island, NY), 50 µM 2-mercaptoethanol (Fisher Scientific, Fair Lawn, NJ), and 1 mg/ml dextrose (Fisher). To this mixture was added human interferon-γ, 20 U/ml, (PeproTech, Inc., Rocky Hill, NJ) and Pansorbin cells (formalinized *Staphylococcus aureus*, Cowan strain; Calbiochem, San Diego, CA) at a final dilution of 1/4000. (Prior to use in the cultures, Pansorbin cells were washed 2 times with Dulbecco's phosphate-buffered saline (GIBCO BRL) and reconstituted to the same volume as supplied by the manufacturer.) The resulting cell suspension was aliquoted into 162 cm² tissue culture flasks (Costar, Cambridge, MA), 80 ml/flask, and the flasks were incubated horizontally at 37°C in a humidified atmosphere of 5% CO₂/95% air for 24 hours. The culture supernatant fluids were then harvested by centrifugation and sterilized by filtration through a 0.22 µm filter (Costar). IL-12 heterodimer plus IL-12 p40 were purified from the culture supernatants by immunoaffinity chromatography using a 2-4A1 protein G sepharose (PGS) column, as described below for the purification of rhesus IL-12, except that the elution buffer contained 0.01% gelatin (Sigma) to minimize protein loss due to nonspecific adsorption to surfaces. The eluate was dialyzed for 4 to 6 hours against 100-200 volumes of Dulbecco's phosphate-buffered saline, and then overnight against the same volume of RPMI 1640 containing 100 µg/ml gentamicin. The dialyzed eluates were sterilized by passage through a 0.22 µm filter, and then assayed by ELISA for content of IL-12 heterodimer and IL-12 p40 (Gately, M.K., Chizzonite, R. and

Presky, D.H., Measurement of human and mouse interleukin 12, *Current Protocols in Immunology*, vol. 1. J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, and W. Strober, eds., John Wiley & Sons, Inc., New York, 1995, pp. 6.16.1-6.16.15) and for IL-12 bioactivity (ibid.). Typically, the weight ratio of IL-12 p40:IL-12 heterodimer, as measured by ELISA, was approximately 5:1.

5

EXAMPLE 2

Production of Recombinant Human IL-12

Recombinant Human IL-12 was prepared, characterized, and generated as set forth in U.S. Patent No. 5,536,657 the contents of which are expressly incorporated herein by reference.

EXAMPLE 3

Generation of Rhesus Monkey IL-12

The p35 and p40 subunit cDNA sequences for rhesus monkey IL-12 (F.Villinger et al., J. Immunol., 155:3946-3954 (1995)) were engineered for expression in CHO-dhfr minus cells on two separate plasmids using standard procedures (Current protocols in molecular biology, F. Ausubel, ed., J. Wiley and Sons, Inc., New York (1993)). Clones were obtained from an unamplified cell population and their IL-12 production was monitored by an IL-12 specific ELISA. An optimally producing clone was selected and adapted to growth in CHO serum-free medium (Sigma). The cells were subsequently grown in spinner cultures for protein production purposes. Rhesus monkey IL-12 was purified from the supernatants by antibody affinity chromatography. The affinity column was produced by crosslinking 10 mg of anti-human IL-12 p40 mAb 2-4A1 (Chizzonite et al., J. Immunol., 147:1548-1556 (1991)) to Protein G Sepharose (Pharmacia Biotech) using 10 mM dimethyl pimelimidate (Pierce, Rockford, IL) at a density of 1 mg mAb/ml of gel (Stern and Podlaski, Tech. In

Protein Chem. IV, Acad. Press, New York, 353-360 (1993)). The serum-free CHO supernatant containing rhesus IL-12 was filtered through a 0.2 μ m filter and loaded directly onto the 10 ml 2-4A1 PGS column previously equilibrated in PBS pH 7.2. The flow rate was 1 ml/min. The column was washed with 10 volumes of PBS and eluted with 0.1M Glycine-HCL, 0.15M NaCl pH 3.0. The eluate was immediately neutralized with 3M Tris-HCl, pH 9. The affinity column was able to bind ~2 mgs of rhesus IL-12/run, including excess p40 monomer, as determined by Bradford and SDS-PAGE. Other contaminants were at trace levels. To concentrate and further purify the rhesus IL-12, the IL-12-containing eluate was dialyzed against 20 mM Na Phosphate, pH 7, and loaded onto a S-Sepharose column conditioned with the same buffer solution. The flow rate was 1 ml/min. All protein was bound. The column was washed with 10 volumes of phosphate buffer and then eluted with phosphate buffer containing 0.3M NaCl. The eluted pool was assayed for endotoxin using the LAL kit from Biowittaker and found to be < 10 EU/mg of protein. Western Blot analysis using the mAb 2-4A1 as detection reagent showed the rhesus IL-12 heterodimer at ~80 kDa as well as an apparent excess of p40 monomer at 40 kDa. Coomassie stained SDS-PAGE shows an additional prominent protein of equal intensity to the p80 heterodimer at ~70 kDa. Both the 80 and 70 kDa proteins are reduced to their monomeric forms after treatment with 2-mercaptoethanol but the latter protein band does not react with mAb 2-4A1.

EXAMPLE 4

Preparation, Characterization, and Purification of Hybridoma Antibodies

5 Mice carrying a mutation in the IL-12 p35 subunit gene on the Balb/c background were produced as described in Mattner, F., et al., *Eur. J. Immunol.*, 26:1553-1559 (1996). The IL-12 p35-deficient mice were immunized intraperitoneally with 5 µg of purified recombinant human IL-12 in complete Freund's adjuvant. The mice received 3 subsequent intraperitoneal booster injections of 5 µg human IL-12 in incomplete Freund's adjuvant over a 2.5 month period. Final injections of 75 µg human IL-12 in PBS (50 µg i.p. and 25 µg i.v.) were given three and two days before splenectomy, followed by an i.p. injection of 50 µg of human IL-12 in PBS one day before splenectomy. Splenocytes were harvested from these mice and fused to mouse myeloma SP2/0 cells at a ratio of 1:1 using 50% w/v polyethylene glycol 1500 (Boehringer Mannheim) according to the method of Oi and Herzenberg, in *Selected Methods in Cellular Immunology*, ed. B. Mishell and S. Shiigi, W. H. Freeman and Co., New York, 1980, pp.351 - 372. The fused cells were plated at a density of 60,000 total cells/well in 96-well cluster plates in IMDM supplemented with 10% FBS (Hyclone), 100 Units/ml penicillin G (BioWhittaker), 100 µg/ml streptomycin (BioWhittaker), 250 ng/ml Fungizone (BioWhittaker), 2 mM glutamine (BioWhittaker), 100 µg/ml gentamicin sulfate (BioWhittaker), 50 µM 2-mercaptoethanol (BioRad), 100 µM hypoxanthine (Sigma), 400 nM aminopterin (Sigma), 16 µM thymidine (Sigma), and 2.5% P388D1 supernatant (produced as described by Nordan, R.P., et al., *J. Immunol.*, 139:813 (1987)). Hybridoma supernatants were assayed for specific anti-human IL-12 antibodies by immunoprecipitation of ¹²⁵I-labeled human IL-12 as described below. Hybridoma cell lines secreting anti-human IL-12 antibodies were cloned by limiting dilution. Antibodies were purified from ascites by sequential treatment with caprylic acid and ammonium sulfate as previously described (Reik, L. et al., *J. Immunol. Methods*, 100:123-130 (1987)).

EXAMPLE 5

Preparation of ^{125}I -labeled Human IL-12

5

Recombinant human IL-12 was radiolabeled to a specific activity of about 2200 Ci/mmol using a modification of the Iodogen (Pierce Chemical Co.) procedure previously described in Chizzonite et al., J. Immunol., 147: 1548-1556 (1991) and Chizzonite et al., J. Immunol., 148: 3117-3124 (1992) which are incorporated herein by reference. Iodogen was dissolved in chloroform and 0.05 mg dried in a 12 x 15 mm borosilicate glass tube. For radiolabeling, 1.0 mCi Na[^{125}I] (Amersham, Chicago, Ill.) was added to an Iodogen-coated tube containing 0.05 ml of Tris-iodination buffer (25 mM Tris-HCL pH 7.5, 0.4 M NaCl and 1mM EDTA) and incubated for 6 min at room temperature. The activated ^{125}I solution was transferred to a tube containing 0.1 ml IL-12 (31.5 μg) in Tris-iodination buffer and the reaction was incubated for 6 minutes at room temperature. At the end of the incubation, 0.05 ml of Iodogen stop buffer (10 mg/ml tyrosine, 10% glycerol in Dulbecco's PBS, pH 7.40) was added and reacted for 5 minutes. The mixture was then diluted with 1% (w/v) BSA in 1.0 ml Tris-iodination buffer, and applied to a Bio-Gel P10DG desalting column (BioRad Laboratories (BRL)) for chromatography. The column was eluted with 1% (w/v) BSA in Tris-iodination buffer, and fractions (1 ml) containing the peak amounts of labeled protein were combined and diluted to 1×10^8 cpm/ml with 1% (w/v) BSA in Tris-iodination buffer. The TCA precipitable radioactivity (10% TCA final concentration) was typically in excess of 95% of the total radioactivity. The radiospecific activity of the recombinant human IL-12 was typically about 2200 Ci/mmol.

20

EXAMPLE 6

Immunoprecipitation Assay of ^{125}I -labeled Human IL-12

5 Nunc Maxisorp 96-well break-apart plates were coated with rabbit affinity purified antibody to mouse IgG (Cappel, Durham, NC) by incubating 18 hrs at 4° C with 100 μl /well of 5 $\mu\text{g}/\text{ml}$ rabbit anti-mouse IgG in carbonate coating buffer (15 mM Na_2CO_3 / 35 mM NaHCO_3), pH 9.6. The coated wells were washed with PBS /0.05% Tween-20/0.01% Thimerosol and then blocked by incubation with 200 μl of 1% (w/v) BSA/PBS/0.01% Thimerosol for 4 hrs at 37° C. Hybridoma supernatants (75 μl) were added to the anti-mouse IgG-coated wells and incubated for 3 hrs at 22° C. The wells were washed 3 times with 300 μl of PBS/0.05% Tween-20/0.01% Thimerosol, and then 100,000 cpm of ^{125}I -labeled human IL-12 were added to each well in 100 μl of antibody diluting buffer (PBS/1% BSA (w/v)/0.5 M NaCl/0.05% Tween-20/0.01% Thimerosol). After 18 hrs at 4° C, the wells were washed 3 times with 200 μl of PBS/0.05% Tween-20/0.01% Thimerosol. The wells were then separated and the amount of radioactivity bound to the wells was determined using a gamma counter. In some experiments, following the incubation of the hybridoma supernatants in the rabbit anti-mouse IgG-coated wells, 100 μl of conditioned supernatant from human IL-12 p40-transfected COS cells prepared as previously described (Gubler et al., Proc. Natl. Acad. Sci. 88: 4143-4147 (1991)) were incubated in the wells for 1 hr at 37° C prior to addition of ^{125}I -labeled human IL-12 to determine if
20 the captured mouse anti-human IL-12 antibodies bound to the p40 subunit of human IL-12.

EXAMPLE 7

Identification of Monoclonal Anti-Human IL-12 Antibodies

5 A 96-well plate-based immunoprecipitation assay was used to identify hybridomas secreting anti-human IL-12 antibodies. Hybridoma supernatants were incubated in the absence and presence of 100µl COS cell supernatant containing human IL-12 p40 subunit as described above. ¹²⁵I-labeled human IL-12 (100,000 cpm/well) was added, and the amount of ¹²⁵I-labeled human IL-12 captured onto the wells was determined. Fig. 1 shows that antibodies contained in supernatants from hybridomas 5F2, 16F2, 16G2, 20E11 and 17E2 captured ¹²⁵I-labeled human IL-12. In addition, the presence of unlabeled human IL-12 p40 subunit during the immunoprecipitation reaction did not block capture of ¹²⁵I-labeled human IL-12 by the antibodies 5F2, 16F2, 16G2 and 20E11, demonstrating that these antibodies do not have high affinity for the IL-12 p40 subunit alone. In contrast, the presence of unlabeled human IL-12 p40 subunit during the immunoprecipitation reaction completely blocked capture of ¹²⁵I-labeled human IL-12 by 17E2, demonstrating that 17E2 recognized the p40 subunit of human IL-12.

EXAMPLE 8

Analytical Isoelectric Focusing of Anti-Human IL-12 Monoclonal Antibodies

Analytical isoelectric focusing was performed using a pH 3.5-9.5 Ampholine PAGplate from Pharmacia Biotech (code no. 80-1124-80, Uppsala, Sweden). Isoelectric focusing was done according to the manufacturer's instructions using electrode solutions of 1 M phosphoric acid and 1N sodium hydroxide. The gel was loaded with 5 samples, each of which contained a single immunoglobulin, i.e., 20E11, 5F2, 20C2, 16G2 and 16F2. Standards were from the Isoelectric Focusing pH 3-10 Calibration Kit from Pharmacia Biotech (code no. 17-0471-01). Running conditions were 1000 volts, 10 watts, 2.5

hours, 4°C. The gel was silver stained using the Pharmacia Biotech PlusOne Silver Staining Kit for protein (code no. 17-1150-01) according to the manufacturer's directions.

Fig. 2 shows the isoelectric focusing patterns of anti-human IL-12 monoclonal antibodies 20C2, 16G2, 16F2, 20E11, and 5F2.

5

EXAMPLE 9

Isoelectric Focusing Patterns of the Anti-Human IL-12 Monoclonal Antibodies

As shown in Fig. 2, monoclonal antibodies 20C2 (Chizzonite et al., Cytokine, 6: A82a (1994)), 20E11, and 5F2 are unique immunoglobulins. Monoclonal antibodies 16G2 and 16F2 appear identical by isoelectric focusing, but both are different from 20C2, 20E11 and 5F2. The pI of these antibodies is in the range of pH 5-6.

EXAMPLE 10

Generation of PHA-Activated Lymphoblasts

Day 4 PHA-activated human peripheral blood mononuclear cells (PBMC) were used in determining both natural human IL-12-induced and rhesus monkey IL-12-induced proliferation. PBMC were isolated (Gately et al., J. Natl. Cancer Inst., 69:1245 (1982)) and stimulated with 0.1% PHA-P (Difco Labs., Detroit, MI). After 3 days, the cultures were split 1:1 with fresh medium and recombinant 50 U/ml human IL-2 (provided by Dr. F. Khan, Hoffmann-La Roche Inc.) as described (Gately, M.K., Chizzonite, R. and Presky, D.H., Measurement of human and mouse interleukin 12, *Current Protocols in Immunology*, vol. 1, J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, and W. Strober, eds., John Wiley & Sons, Inc., New York, 1995, pp. 6.16.1-6.16.15).

Supplemented medium used for cell culture was as described previously for the production of natural human IL-12 with the addition of 5% human AB serum (Irvine Scientific, Santa Ana, CA).

EXAMPLE 11

5

Lymphocyte Proliferation Assay

The effects of the various anti-human IL-12 monoclonal antibodies on IL-12- and IL-2-stimulated PHA-activated human lymphoblast proliferation was determined by a method based on M. K. Gately et al. (Gately, M.K., Chizzonite, R. and Presky, D.H., Measurement of human and mouse interleukin 12, *Current Protocols in Immunology*, vol. 1., J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, and W. Strober, eds., John Wiley & Sons, Inc., New York, 1995, pp. 6.16.1-6.16.15). Day 4 PHA-activated lymphoblasts, prepared as described above, were harvested, washed and resuspended in supplemented medium at 4×10^5 cells/ml and incubated in 96-well plates (2×10^4 cells/well) with purified monoclonal anti-human IL-12 antibody and the relevant cytokine, i.e. human or monkey IL-12. Twenty five μ l aliquots of both natural human IL-12 at 1 ng/ml or monkey IL-12 at 2 ng/ml were mixed with 25 μ l aliquots of various dilutions of anti-human IL-12 monoclonal antibodies (mAbs). The final antibody concentration in the wells varied from 0.0005 μ g/ml up to 0.5 μ g/ml. A separate, identical set of wells containing the various anti-human IL-12 mAbs and recombinant IL-2 was prepared to determine the effects of the anti-human IL-12 mAbs on IL-2-stimulated proliferation as a measure of inhibitory specificity. A standard dose-response curve ranging from 250 pg or 500 pg per well human or monkey IL-12, respectively, down to 0 pg without added antibodies was also included to determine IL-12-responsiveness. Plates containing mixtures of cytokines and antibodies were incubated for 30 minutes at 37°C, and then 50 μ l aliquots of cell suspension were added to the wells. The culture plates were maintained at 37°C in a humidified atmosphere of 5% CO₂ in air for 48 hours prior to ³H-thymidine pulsing. Fifty μ l of 10 μ Ci/ml ³H-thymidine (diluted in supplemented media with 5% FCS in lieu of 5% human AB serum) were

added to each well. After incubation for an additional 6 hr at 37° C, the well contents were harvested onto glass fiber filters via a cell harvester, and ³H-thymidine incorporation into cellular DNA was measured by use of a liquid scintillation counter. Values shown in Figs. 3 and 4 are the means of triplicate wells.

5

EXAMPLE 12

Inhibition of Cytokine-Stimulated PHA-Activated Lymphoblast Proliferation by Monoclonal Anti-Human IL-12 Antibodies

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

Proliferation of PHA-activated human lymphoblasts stimulated with 0.25 ng/ml IL-12 was inhibited in a dose-dependent fashion by antibodies 5F2, 16F2, 16G2, and 20E11 (Fig. 3). The potencies of these anti-human antibodies, defined as the concentration that produces 50% of maximal inhibition (IC₅₀) of 0.25 ng/ml IL-12-stimulated proliferation, are 0.03 µg/ml for 5F2, 0.01 µg/ml for 16F2, 0.01 µg/ml for 16G2, and 0.01 µg/ml for 20E11. The maximal (9440 cpm) and background (1480 cpm) levels of lymphoblast proliferation are represented by the horizontal dotted lines at the upper and lower ends of the plots, respectively. As shown in Fig. 3, the 5F2, 16F2, 16G2, and 20E11 antibodies were able to inhibit human IL-12 stimulated PHA activated lymphoblast proliferation by at least 90%. In contrast, as also shown in Fig. 3, the previously identified anti-human IL-12 p75-specific antibody 20C2 (Chizzonite et al., Cytokine, 6: A82a (1994)) is not able to substantially inhibit human IL-12 bioactivity.

In addition, as shown in Fig. 4, 5F2, 16F2, 16G2, and 20E11 potently inhibited proliferation of PHA-activated human lymphoblasts stimulated with 0.5 ng/ml rhesus monkey IL-12, with a similar IC₅₀ to that seen with human IL-12-stimulated proliferation. In contrast, 20C2 has only a minimal inhibitory effect on rhesus monkey IL-12-stimulated proliferation. Therefore, the antibodies 5F2, 16F2, 16G2, and 20E11 appear to exhibit good cross-reactivity to rhesus monkey IL-12, whereas the cross-

reactivity of 20C2 is much less. None of these monoclonal antibodies inhibited IL-2-induced proliferation, demonstrating that their effect on IL-12-stimulated proliferation was specific for IL-12 and was not due to a general inhibition of cell proliferation.

5

EXAMPLE 13

Interferon- γ Synthesis Assay

Interferon- γ (IFN- γ) synthesis was induced using Day 4 PHA-activated human lymphoblasts produced as described above. The medium used was a 1:1 mixture of RPMI 1640 and Dulbecco's modified Eagle's medium supplemented as described above for preparation of natural human IL-12 and containing, in addition, 5% heat-inactivated (56°C, 30 min) fetal bovine serum (Hyclone, Logan, UT) in lieu of human AB serum. Duplicate 1 ml cultures were set up in the wells of 24-well tissue culture plates (Costar). To each well was added 5×10^5 PHA-activated lymphoblasts, 0.25 ng/ml purified natural human IL-12, 20 units/ml recombinant human IL-2, 1 ng/ml recombinant human IL-1 β (provided by Dr. R. Chizzonite, Hoffmann-La Roche), and the indicated concentrations of anti-human IL-12 antibodies. Initially, all reagents except the lymphoblasts were added to the wells and incubated at 37°C for 30 min, followed by the addition of the lymphoblasts. The cultures were then incubated for 24 hr at 37°C in a humidified atmosphere of 5% CO₂ in air. At the end of this time, the culture supernatant fluids were harvested by centrifugation and assayed for their content of IFN- γ by use of an ELISA. The amount of IFN- γ produced in cultures containing lymphoblasts with IL-2 + IL-1 but no IL-12 was always less than 15% and usually less than 5% of that produced in cultures containing 0.25 ng/ml IL-12 in addition to IL-2 + IL-1.

25 The ELISA for measuring human IFN- γ used monoclonal anti-human IFN- γ antibodies from Endogen (Woburn, MA). Nunc EIA plates (Fisher) were coated overnight at 4°C with 100 μ l/well of 1 μ g/ml anti-human IFN- γ (Endogen #M-700A) in coating buffer (0.015 M Na₂CO₃ + 0.035 M

NaHCO₃ in distilled water, pH 9.6). The following morning, the coating buffer was flicked out of the wells, and the wells were blocked by addition of 200 µl/well of Dulbecco's phosphate-buffered saline (D-PBS; Fisher) containing 1% bovine serum albumin (Sigma). After incubation for 1 hr at room temperature, the plates were washed with D-PBS containing 0.05% tween 20 (Sigma), and 100 µl
5 aliquots of recombinant human IFN-γ standard (Endogen) or culture supernatants diluted in assay buffer (D-PBS + 0.5% bovine serum albumin + 0.05% tween 20) were added to the wells. The plates were then incubated for 2 hours at room temperature with shaking. Following this, the plates were again washed, and each well received 100 µl of 300 ng/ml biotinylated anti-human IFN-γ (Endogen #M-701-B) in assay buffer. The plates were incubated for 1 hr at 37°C, followed by washing. One hundred µl aliquots of streptavidin-peroxidase (Sigma) diluted 1:1000 in assay buffer were then added to each well, and the plates were incubated for 30 min at 37°C. The plates were again washed and then developed by addition of 100 µl aliquots of a 1:1 mix of TMB Peroxidase Substrate and Peroxidase B Solution (Kirkegaard & Perry Laboratories, Gaithersburg, MD). The reaction was stopped after ~12 min by addition of 50 µl/well of 1 M H₃PO₄, and the absorbance was read at 450 nm with subtraction of background at 650 nm.

EXAMPLE 14

Inhibition of Cytokine-Stimulated Interferon-γ Production by Monoclonal Anti-Human IL-12

Antibodies

Production of IFN-γ by PHA-activated human lymphoblasts stimulated with 0.25 ng/ml human IL-12 was inhibited in a dose-dependent fashion by antibodies 5F2, 16F2, 16G2, and 20E11 (Fig. 5). The potencies of these anti-human antibodies, defined as the concentration that produces 50% of
25 maximal inhibition (IC₅₀) of 0.25 ng/ml human IL-12-stimulated IFN-γ production, are 0.02 µg/ml for 5F2, 0.02 µg/ml for 16F2, 0.01 µg/ml for 16G2, and 0.02 µg/ml for 20E11. These anti-human heterodimer specific IL-12 antibodies were able to inhibit greater than 90% of IL-12-stimulated IFN-γ

production when used at 0.5 µg/ml. In contrast, the previously identified anti-human IL-12 p75-specific antibody 20C2 (Chizzonite et al., Cytokine, 6: A82a (1994)) is less potent and is unable to inhibit IL-12-stimulated IFN-γ production by more than 65% at concentrations less than or equal to 0.5 µg/ml.

5

EXAMPLE 15

Sequence Analysis of the Genes Encoding the Variable Region of the Antibody Heavy Chains Present in the Anti-Human IL-12 Antibody-Producing Hybridoma Cell Lines

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201

EXAMPLE 16

Nucleotide and Deduced Amino Acid Sequences of the Variable Region of the Monoclonal Anti-Human IL-12 Antibody Heavy Chains

5

INSCS)

The nucleotide sequences of a portion of the variable region of the immunoglobulin heavy chain gene encompassing framework region (FR) 1, complementarity determining region (CDR) 1, FR2, CDR2, FR3, CDR3, and FR4 of IL-12 antibodies produced by hybridoma cell lines HIL-12F3-16G2 and HIL-12F3-20E11 and the deduced amino acid sequences thereof are shown in Fig. 6 and Fig. 7, respectively. The CDR sequences are underlined. Comparison of available sequence information showed that the heavy chains of antibodies produced by hybridomas HIL-12F3-16G2 and HIL-12F3-20E11 exhibit 94% homology at the DNA level and 93% similarity at the amino acid level.

10
20
30
40
50
60
70
80
90
100
110
120
130
140
150
160
170
180
190
200
210
220
230
240
250
260
270
280
290
300
310
320
330
340
350
360
370
380
390
400
410
420
430
440
450
460
470
480
490
500
510
520
530
540
550
560
570
580
590
600
610
620
630
640
650
660
670
680
690
700
710
720
730
740
750
760
770
780
790
800
810
820
830
840
850
860
870
880
890
900
910
920
930
940
950
960
970
980
990
1000